

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/322939952>

# Tumor necrosis factor- $\alpha$ 308 G/A polymorphism and type 2 diabetes mellitus; a systematic review and meta-analysis

Article · January 2018

DOI: 10.15171/jnp.2018.09

CITATIONS

0

READS

77

6 authors, including:



**Ali Hassanpour Dehkordi**

Shahrekord University of Medical Sciences

61 PUBLICATIONS 448 CITATIONS

[SEE PROFILE](#)



**Majid Dousti**

Shiraz University of Medical Sciences

8 PUBLICATIONS 47 CITATIONS

[SEE PROFILE](#)



**Mona Zamanian Azodi**

National Elites Foundation

63 PUBLICATIONS 375 CITATIONS

[SEE PROFILE](#)



**Mostafa Rezaei tavirani**

Shahid Beheshti University of Medical Sciences

254 PUBLICATIONS 1,674 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Breast cancer research in Iran: a scientometric analysis of publications output from 1991 to 2015 in Scopus [View project](#)



effectiveness of Furazolidone and Clarithromycin [View project](#)

# Journal of Nephropathology



Click for updates

## Tumor necrosis factor-alpha 308 G/A polymorphism and type 2 diabetes mellitus; a systematic review and meta-analysis

Ali Hasanpour Dehkordi<sup>1,2</sup>, Majid Dousti<sup>3</sup>, Faezeh Kiani<sup>3</sup>, Mona Zamanian Azodi<sup>4</sup>, Mostafa Rezaei-Tavirani<sup>4</sup>, Kourosh Sayehmiri<sup>5</sup>

<sup>1</sup>Social determinants of Health Research Center, Shahrekord University of Medical sciences, Shahrekord, Iran

<sup>2</sup>Department of Medical-Surgical, Faculty of Nursing and Midwifery, Shahrekord University of Medical sciences, Shahrekord, Iran

<sup>3</sup>Student Research Committee, Ilam University of Medical Sciences, Ilam, Iran

<sup>4</sup>Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>5</sup>Department of Biostatistics, Psychosocial Injury Research Center, Ilam University of Medical Sciences, Ilam, Iran

### ARTICLE INFO

*Article type:*  
Review

*Article history:*  
Received: 15 July 2017  
Accepted: 17 December 2017  
Published online: 5 January 2018  
DOI: 10.15171/jnp.2018.09

*Keywords:*  
Diabetes mellitus type 2  
TNF- $\alpha$   
308 G/A  
Polymorphism

### ABSTRACT

**Context:** Type 2 diabetes mellitus (T2DM) is a multi-factorial disease with several genetic and environmental factors known to be associated with the disease. Tumor necrosis factor-alpha (TNF- $\alpha$ ) acts on fat cells associated with increased insulin resistance and obesity and its increase can be associated with the prevalence of T2DM. Previous studies on the relationship between 308 G/A polymorphism in TNF- $\alpha$  and T2DM were not conclusive. The aim of this study was to investigate the association between 308 G/A polymorphism in TNF- $\alpha$  gene and T2DM using a systematic review and meta-analysis.

**Evidence Acquisitions:** In this systematic review we searched all published studies about the association between TNF 308 G/A polymorphism and T2DM through databases such as Scopus, Science direct, PubMed and Google scholar. A fixed or random effect model was used on the basis of heterogeneity. The heterogeneity was assessed using the  $I^2$  index. We used STATA software (version 11.2) for data analysis.

**Results:** In 22 selected papers, the total number of T2DM and control subject was 8485 and 8615, respectively. Odds ratios (ORs) with 95% confidence intervals (CI) for the GG genotype polymorphism 308 G/A in total, Asian populations and whites (Caucasian) were estimated as OR = 0.98 (95% CI: 0.96-1.00), OR = 0.96 (95% CI: 0.93 to 0.99) and OR = 1.00 (95% CI: 0.97 to 1.40). The Publication Bias/or heterogeneity was observed in this study.

**Conclusions:** The present investigation have shown that 308 G/A polymorphism GG and GA genotypes in TNF- $\alpha$  gene cannot be considered as predictors of T2DM ( $P < 0.05$ ).

### Implication for health policy/practice/research/medical education:

In a systematic review on 22 selected papers, we found that 308 G/A polymorphism GG and GA genotypes in TNF- $\alpha$  gene cannot be considered as predictors of type 2 diabetes mellitus ( $P < 0.05$ ).

**Please cite this paper as:** Hasanpour Dehkordi A, Dousti M, Kiani F, Zamanian Azodi M, Rezaei-Tavirani M, Sayehmiri K. Tumor necrosis factor-alpha 308 G/A polymorphism and type 2 diabetes mellitus; a systematic review and meta-analysis. J Nephropathol. 2018;7(1):25-31. DOI: 10.15171/jnp.2018.09.

### 1. Context

Diabetes mellitus (DM) is a chronic disease and is one of the health challenges in the 21st century. Decrease in insulin secretion and/or resistance to it are known to be involved in the development of DM (1). In general, DM refers to a heterogeneous group of metabolic disorders characterized by decreased insulin action or resistance

to it, or both and is divided into two main categories; diabetes type 1, and type 2. DM is generally characteristic by insulin resistance and decrease in insulin production (2). Approximately 90% of the global epidemics of DM belongs to type 2 DM (T2DM) and it has become the fourth common cause of death in most developed countries. According to the international diabetes

\*Corresponding author: Mona Zamanian Azodi, Email: mona.azodi@gmail.com

federation in 2003, the proportion of diabetes, in the world, was 194 million people and it is estimated this number will increase to 366 million people in 2030 which will be mostly T2DM. Thus, due to increased demand for health care and the imposition of economic pressures, diabetes (more specifically T2DM) has become an important challenge for different communities (3).

There are known factors predisposing individuals to T2DM. It is a multi-factorial disease and several factors both genetics and environmental play role in its development. Because of the magnitude, complexity and multi-factorial nature of diabetes, the accurate diagnosis of the genetic factors is difficult. Gene polymorphisms have been considered to be essential for the development of disease. Genomic studies in several countries and different races have been performed to identify genes associated with T2DM (4-6). However, the exact gene (/or genes) involved in T2DM is not recognized. It seems that all genes involved in the pathway of insulin biosynthesis in pancreatic insulin and obesity can be the potential candidates (7). So far, more than 200 genes have been identified. One of these genes is tumor necrosis factor-alpha (TNF- $\alpha$ ). As a regulator, TNF- $\alpha$  acts in fat cells and is associated with increased insulin resistance and obesity (8). Adipose tissue is an important source of TNF- $\alpha$ . Thus TNF- $\alpha$  expression in human adipose tissue and muscle increases the risk of obesity (9,10). One of the known polymorphisms in tumor necrosis factor-alpha (TNF- $\alpha$ ) gene that is located on the short arm of chromosome 6, is the displacement of guanine with adenine at position 308 promotes the mentioned gene. In general, the results of studies on the association between the TNF- $\alpha$  gene polymorphism and T2DM is inconsistent because in different populations different results have been found (11).

Today's chronic diseases are major challenges for governments, nurses, healthcare professionals, families and the community (12-17). So, many social and economic costs are imposed annually on the family and society (18,19).

Despite many studies conducted in this field, in different countries and races, the existence or lack of relationship between 308 G/A polymorphism of TNF- $\alpha$  and the risk of T2DM has not been clearly demonstrated. The aim of this study was to investigate the association of 308 G/A polymorphism in TNF- $\alpha$  gene and T2DM using a large-scale meta-analysis.

## 2. Evidence Acquisitions

### 2.1. Literature search and study selection

To find related documents, we searched the following

literature databases: Scopus, PubMed, and Google scholar, SID, Irandoc and Magiran. We performed article search mainly using systematic search for all possible combinations of English key words, the major and sensitive ones and their equivalent keywords. We searched based on keywords; TNF- $\alpha$ , type 2 DM and polymorphism. To determine the scope of this study, case-control studies and meta-analysis of previous articles were collected. The article selection was as follows; 1) a related case-control study and 2) having complete data about genotype and adequate alleles. All searches were performed independently by two people. Under this process, in initial inquiry we found 37 articles conducted during 1995 to 2013. The main criterion to select papers in this study, was referring to the association between gene, polymorphisms of TNF- $\alpha$  with T2DM. We added all relevant papers to the initial list. Studies about clinical decision making or those about clinical characteristics of patients or those not related to the topic of the research are excluded. The 8 non-relevant articles were removed and 29 articles' abstracts were investigated. Due to the unavailability of full texts, the 2 papers' abstracts were deleted and the full texts of 27 articles were investigated for the quality and finally, 22 articles were analyzed (Figure 1).

### 2.2. Statistical analysis

We extracted from each article the following data; the number of controls, number of cases, the presence of genotypes GG, GA and AA polymorphism 308 G/A in cases and controls, age, sex and year of publication




Figure 1. Study flowchart.

of the paper. Odds ratios (ORs) and its 95% confidence intervals (CIs) were calculated for each study.

Peto method was used to calculate OR and pooled OR. The heterogeneity was assessed using the Q and  $I^2$  indexes. Begg's funnel plot was used to check publication bias. The data were analyzed using R program (version 2/15/1) and Stata (version 11.2). Additionally  $P$  value less than 0.05 was considered as statistical significance.

### 3. Results

In 22 reviewed papers, the total number of T2DM and control subjects were 8485 and 8615, including 6003 cases and 5683 controls with genotype GG, 2136 cases and 2433 controls with GA. AA genotypes were found among 198 cases and 408 controls. The European, Australian, Mexican, Japanese, Indian, Chinese, Brazilian, Chilean, and Iranian populations have been studied from 1995 to 2013. Characteristics of the studies are summarized in Table 1.

According to the results from the Random Effects Model, OR (0.99) and 95% CIs (from 0.97 to 1.01) were estimated. There was no significant association between GG genotype and T2DM ( $P > 0.05$ ). Due to the heterogeneity of the studies ( $I^2$  Index=1.82), the CIs for

each study based on a Random Effects Model is given in Figure 1.

The results of three studies carried out by Zeggini et al,<sup>27</sup> Guzmán-Flores et al,<sup>22</sup> and Golshani et al, to GG genotype illustrate its protective role against T2DM (Figure 1), while in study conducted by Boraska et al,<sup>26</sup> it has been indicated that individuals having GG genotype are 1.46-folds higher at risk of T2DM. In the 17 studies, there was no significant difference between GG genotype and the susceptibility to T2DM. Combining the results of all 21 studies, it has been shown no significant relationship between GG genotype polymorphism 308 G/A genes in TNF- $\alpha$  and susceptibility to T2DM (Figure 1).


In Figure 2, we found the publication bias has no influence on GG genotype indicated by the symmetry of the funnel. The size of the circles has shown the precision of the studies. Thus, the greater the circles, the greater the samples sizes, and vice versa (Figure 3).

### 4. Discussion

The results of the present study have shown that 308 G/A polymorphism GG and GA genotypes in TNF- $\alpha$  gene cannot be considered as predictors of T2DM

**Table 1.** The characteristics of different articles related to 308G/A (TNF- $\alpha$ ) and susceptibility to type 2 diabetes mellitus

Authors' name	Publication year	Country	Case (n)	Control (n)	Case/Control (age year)	GG		GA+ AA		References
						Case	Total	Case	Total	
Heijmans et al	2002	Netherlands	79	577	>85	51	429	28	227	(20)
Perez-Luque et al	2012	Mexico	95	135	49.5/52	72	188	23	42	(21)
Guzmán-Flores et al	2011	Mexico	518	1290	—	438	1581	80	227	(22)
Li et al	2003	Finland	395	284	65.7/55.0	273	462	122	217	(23)
Furuta et al	2002	Japan	132	142	55.6/51.8	129	268	3	6	(24)
Bouhaha et al	2010	Tunis	228	300	43.8/60.6	204	345	95	149	(25)
Boraska et al	2010	UK	1454	2504	—	938	2571	516	1387	(26)
Zeggini et al	2005	UK	834	1231	—	484	1263	292	726	(27)
Kim et al	2006	Korea	169	198	—	174	315	24	52	(28)
Lindholm et al	2008	Sweden	2957	206	61.2/59.7	1908	2041	1019	1091	(29)
Ishii et al	2000	Japan	71	299	58.76/41.96	69	359	2	11	(30)
Morris et al	2003	Australia	91	189	—	52	178	38	101	(31)
Liu et al	2008	China	245	122	48.2/46.5	222	331	23	36	(32)
Hamann et al	1995	German	138	57	57.9/56.1	108	154	30	41	(33)
Padovani et al	2000	Brazil	21	145	43/42	17	128	4	38	(34)
Ko et al	2003	China	339	202	38.2/36.8	284	455	55	86	(35)
Santos et al	2006	Chile	30	53	69–60	27	72	3	11	(26)
Shiau et al	2003	China	261	189	58.73/58.16	218	386	39	58	(37)
Tsiavou et al	2004	Greece	32	39	51/44	29	61	3	10	(38)
Vendrell et al	2003	Spain	135	207	56.67	109	268	26	74	(39)
Dutta et al	2013	India	122	100	—	76	147	46	75	(40)
Golshani et al	2013	Iran	173	173	50.4/54.1	146	308	27	38	Unpublished
Total			8485	8615		6003	12310	2498	4703	



**Figure 2.** Meta-analysis of the GG genotype 308G/A with T2DM. Square represent effect estimate of individual studies with their 95% confidence intervals with size of squares proportional to the weight assigned to the study in the meta-analysis


( $P < 0.05$ ).

The results of a similar study conducted by Feng et al in 2010 (41) have shown no significant relationship between gene 308 G/A polymorphism TNF- $\alpha$  and T2DM.

In this study we added the result of four conducted studies from 2010 to 2013 to the result of study performed by Feng et al. In his another work, Feng shows that the high levels of polymorphism 308 G/A allele of TNF- $\alpha$  gene is a risk factor for diabetes type 1 (42). In a research by Furuta et al carried in Japan on 308 G/A, 132 diabetic patients and 142 healthy controls, the association between single nucleotide polymorphisms in the TNF- $\alpha$  gene promoter region was found in diabetic patients and controls (24).

Mukhopadhyaya et al, in a work performed by the Indian population of 40 diabetic patients and 40 control subjects have found an association between single nucleotide polymorphisms in the promoter region of genes 308-G/A TNF- $\alpha$  and T2DM (43).

Costa et al evaluated 36 patients with diabetes and 41 control subjects. Low 863 G/A TNF- $\alpha$  allele and a high concentration of soluble receptor-2 (TNFR2) TNF- $\alpha$  were associated with the higher incidence of diabetes type 2. This article has shown that there is an abnormal increase of TNF- $\alpha$  disease, which is closely related to genetics, metabolism and indicated T2DM indexes (44). This meta-analysis study aimed to investigate the association between polymorphisms in G/A 308 gene TNF- $\alpha$  and metabolic syndrome. The findings demonstrate that allele A-TNF increases the risk of metabolic syndrome (45). Additionally, in another study, a significant association between increased body mass



**Figure 3.** Begg's Funnel plot for evaluation of publication bias.

index and polymorphisms has been reported in Swedish women (29).

In a meta-analysis conducted on Han Chinese population, 10 studies including 1425 diabetic patients and 1116 healthy controls were studied. The results have shown that A<G-308 polymorphism is a major risk factor for T2DM and at the locus of A allele may be critical for the development of T2DM in the Han Chinese population (46). Results have shown that in other races, including the Whites (27,28,33) and Japanese (24,30) polymorphism A<G 308 TNF- $\alpha$  is not a risk factor for the development of T2DM. Hence, it is concluded that the race of the subjects is an important factor in making final conclusion.

According to previous studies conducted in different populations, the overall conclusion of this study was that there was no significant association between allelic sequences in 308 G/A gene for tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and the risk of T2DM. However, T2DM and insulin resistance is multi-factorial disorders in which diverse genetic locus susceptibility involved in disease occurrence. Environmental factors (such as nutrition and physical activity) are also, effective in development of this phenotypic occurrence (47). Besides, other factors such as obesity, lack of physical activity and diet are involved in the development of such disease (48). In general, the result of TNF- $\alpha$  gene polymorphism and T2DM is inconsistent. Depending on the population studied, it shows various effects (11). Here, we should remind that populations with a high incidence of T2DM are widespread, for example in the Pima Indian population that cannot be dedicated to environmental factors (49).

## 5. Conclusions

In general, it is suggested that we need to have in future



research, not only to recruit more samples but also focus on environmental factors such as ethnic differences.

#### Authors' contribution

AHD and MZA; The concept, design, data analysis, and manuscript preparation and final revision. MD, FK and MRT; performing search, data collection, data extraction and writing proposal. AHD and KS; statistical analysis, manuscript editing, and manuscript review. MZA; data collection and providing first draft and submission.

#### Conflicts of interest

The authors declare that they have no conflict of interest regarding the contents of this article.

#### Ethical considerations

Ethical issues (including plagiarism, data fabrication, double publication) have been completely observed by the authors.

#### Funding/Support

None.

#### References

- Matthaei S, Stumvoll M, Kellerer M, Haring HU. Pathophysiology and pharmacological treatment of insulin resistance. *Endocr Rev.* 2007;21(6):585-618. doi: 10.1210/edrv.21.6.0413.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2004; 27(1):5-10. doi: 10.2337/diacare.27.2007.S5.
- Meshkani R, Taghikhani M, Mosapour A. 1484insG polymorphism of the *PTPN1* gene is associated with insulin resistance in an Iranian population. *Arch Med Res.* 2007;38(5):556-62. doi: 10.1016/j.arcmed.2007.01.010.
- Bhattacharya S, Dey D, Roy SS. Molecular mechanism of insulin resistance. *J Biosci.* 2007;32(2):556-62.
- Grarup N, Urhammer S, Ek J. Studies of the relationship between the ENPP1 K121Q polymorphism and type 2 diabetes, insulin resistance and obesity in 7333 Danish white subjects. *Diabetologia.* 2006;49(9):2097-104.
- Wiltshire S, Hattersley AT, Hitman GA. A genome-wide scan for loci predisposing to type 2 diabetes in a UK population (the Diabetes UK Warren 2 Repository): Analysis of 573 pedigrees provides independent replication of a susceptibility locus on chromosome 1q. *Am J Hum Genet.* 2001;69(3):553-69. doi: 10.1086/323249.
- Meshkani R, Taghikhani M, Alkateb H. Polymorphisms within the protein tyrosine phosphatase IB (*PTPN1*) gene promoter: Functional characterization and association with type 2 diabetes and related metabolic traits. *Clin Chem.* 2007;53(9):1585-92.
- Hofmann C, Lorenz K. Altered gene expression for tumor necrosis factor- $\alpha$  and its receptors during drug and dietary modulation of insulin resistance. *Endocrinology.* 1994; 134(1):264-70. doi: 10.1210/endo.134.1.8275942.
- Saghizadeh M, Ong JM, Garvey WT, Henry RR, Kern PA. The expression of TNF  $\alpha$  by human muscle. Relationship to insulin resistance. *J Clin Invest.* 1996; 97(4):1111. doi: 10.1172/JCI118504
- Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- $\alpha$ : direct role in obesity-linked insulin resistance. *Science.* 1993;259(5091):87-91.
- Hoffstedt J, Eriksson P, Hellstrom L, Rossner S, Ryden M, Arner P. Excessive fat accumulation is associated with the TNF-  $\alpha$ -308 G/A promoter polymorphism in women but not in men. *Diabetologia.* 2000;43(1):117-20.
- Hasanpour-Dehkordi A. Self-care concept analysis in cancer patients: an evolutionary concept analysis. *Indian J Palliat Care.* 2016;22(4):388. doi: 10.4103/0973-1075.191753.
- Dehkordi AH, Heydarnejad MS. Effect of booklet and combined method on parents' awareness of children with beta-thalassemia major disorder. *J Pak Med Assoc.* 2008;58(9):485-7
- Hasanpour-Dehkordi A, Khaledi-Far A, Khaledi-Far B, Salehi-Tali S. The effect of family training and support on the quality of life and cost of hospital readmissions in congestive heart failure patients in Iran. *Appl Nurs Res.* 2016;31:165-9. doi: 10.1016/j.apnr.2016.03.005.
- Hassanpour Dehkordi A, Mohammadi N, Nikbakht-Nasrabadi A. Hepatitis-related stigma in chronic patients: A qualitative study. *Appl Nurs Res.* 2016;29:206-10. doi: 10.1016/j.apnr.2015.04.010.
- Noghabi AAA, Zandi M, Mehran A, Alavian SM, Dehkordi AH. The effect of education on quality of life in patients under interferon therapy. *Hepat Mon.* 2010;10(3):218-22.
- Shahbazi K, Solati K, Hasanpour-Dehkordi A. Comparison of hypnotherapy and standard medical treatment alone on quality of life in patients with irritable bowel syndrome: a randomized control trial. *J Clin Diagn Res.* 2016;10(5):OC01-4. doi: 10.7860/JCDR/2016/17631.7713.
- Hasanpour-Dehkordi A, Jivad N, Solati K. Effects of yoga on physiological indices, anxiety and social functioning in multiple sclerosis patients: a randomized trial. *J Clin Diagn Res.* 2016;10(6):VC01-5. doi: 10.7860/JCDR/2016/18204.7916.
- Hasanpour-Dehkordi A, Mohammadi N, Nikbakht-Nasrabadi A. Re-designing Orem's self-care theory for patients with chronic hepatitis. *Indian Indian J Palliat Care.* 2016;22(4):395-401. doi: 10.4103/0973-1075.191754 .
- Heijmans BT, Westendorp RG, Droog S, Kluft C, Knook DL. Association of the tumour necrosis factor  $\alpha$  -308G/A polymorphism with the risk of diabetes in an elderly population-based cohort. *Genes Immun.* 2002;

- 3(4):225-8. doi:10.1038/sj.gene.6363859.
21. Perez-Luque E, Malacara JM, Garay-Sevilla ME, Fajardo ME. Association of the TNF- $\alpha$  -308G/A polymorphism with family history of type 2 diabetes mellitus in a Mexican population. *Clin Biochem*. 2012; 45(1):12-5. doi: 10.1016/j.clinbiochem.2011.09.018.
22. Guzmán-Flores JM, Muñoz-Valle JF, Sánchez-Corona J. Tumor necrosis factor-alpha gene promoter -308G/A and -238G/A polymorphisms in Mexican patients with type 2 diabetes mellitus. *Dis Markers*. 2011; 30(1):19-24. doi: 10.3233/DMA-2011-0759.
23. Li H, Groop L, Nilsson A, Weng J, Tuomi TA. Combination of human leukocyte antigen DQB1\*02 and the tumor necrosis factor alpha promoter G308A polymorphism predisposes to an insulin-deficient phenotype in patients with type 2 diabetes. *J Clin Endocrinol Metab*. 2003;88(6):2767-74. doi: org/10.1210/jc.2002-020506.
24. Furuta M, Yano Y, Ito K, Gabazza EC, Katsuki A. Relationship of the tumor necrosis factor-alpha 2308 A/G promoter polymorphism with insulin sensitivity and abdominal fat distribution in Japanese patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract*. 2002;56(2):141-5. doi: 10.1016/S0168-8227(01)00358-8.
25. Bouhaha R, Baroudi T, Ennafta H, Vaillant E, Abid H. Study of TNFalpha 2308G/A and IL6 2174G/C polymorphisms in type 2 diabetes and obesity risk in the Tunisian population. *Clin Biochem*. 2010;43(6):549-52. doi: 10.1016/j.clinbiochem.2010.01.008.
26. Boraska V, Rayner NW, Groves CJ, Frayling TM, Diakite M. Large-scale association analysis of TNF/LTA gene region polymorphisms in type 2 diabetes. *BMC Med Genet*. 2010;11(1):69. doi: 10.1186/1471-2350-11-69.
27. Zeggini E, Groves CJ, Parkinson JR, Halford S, Owen KR. Large scale studies of the association between variation at the TNF/LTA locus and susceptibility to type 2 diabetes. *Diabetologia*. 2005;48(10):2013-7.
28. Kim HR, Lee MK, Park AJ. The 2308 and 2238 polymorphisms of the TNF-alpha promoter gene in type 2 diabetes mellitus. *Korean J Lab Med*. 2006;26(1):58-63. doi: 10.3343/kjlm.2006.26.1.58
29. Lindholm E, Bakhtadze E, Cilio C, Agardh E, Groop L. Association between LTA, TNF and AGER polymorphisms and late diabetic complications. *PLoS one*. 2008;3(6):e2546. doi: 10.1371/journal.pone.0002546.
30. Ishii T, Hirose H, Saito I, Nishikai K, Maruyama H. Tumor necrosis factor alpha gene G-308A polymorphism, insulin resistance, and fasting plasma glucose in young, older, and diabetic Japanese men. *Metabolism*. 2000;49(12):1616-8. doi:10.1053/meta.2000.18560.
31. Morris AM, Heilbronn LK, Noakes M, Kind KL, Clifton PM. 2308 Nco I polymorphism of tumour necrosis factor alpha in overweight Caucasians. *Diabetes Res Clin Pract*. 2003;62(3):197-201. doi: 10.1016/j.diabres.2003.08.002.
32. Liu HL, Lin YG, Wu J, Sun H, Gong ZC. Impact of genetic polymorphisms of leptin and TNF-alpha on rosiglitazone response in Chinese patients with type 2 diabetes. *Eur J Clin Pharmacol*. 2008;64(7):663-71.
33. Hamann A, Mantzoros C, Vidal-Puig A, Flier JS. Genetic variability in the TNF-alpha promoter is not associated with type II diabetes mellitus (NIDDM). *Biochem Biophys Res Commun*. 1995; 211(3):833-9. doi: 10.1006/bbrc.1995.1887
34. Padovani JC, Pazin-Filho A, Simoes MV, Marin-Neto JA, Zago MA. Gene polymorphisms in the TNF locus and the risk of myocardial infarction. *Thromb Res*. 2000; 100(4):263-9. doi: 10.1016/S0049-3848(00)00315-7.
35. Ko GT, Lee SC, Pu YB, Ng MC, So WY. Tumour necrosis factor alpha promoter gene polymorphism at 2308 (genotype AA) in Chinese subjects with Type 2 diabetes. *Diabet Med*. 2000;20(2):167-8. doi: 10.1046/j.1464-5491.2003.00829\_1.x.
36. Santos MJ, Patino GA, Angel BB, Martinez HJ, Perez BF. Association between tumor necrosis factor-alpha promoter polymorphisms and type 2 diabetes and obesity in Chilean elderly women. *Rev Med Chil*. 2006;134(9):1099-106. doi: 10.4067/S0034-98872006000900003.
37. Shiau MY, Wu CY, Huang CN, Hu SW, Lin SJ. TNF-alpha polymorphisms and type 2 diabetes mellitus in Taiwanese patients. *Tissue Antigens*. 2003;61(5):393-7. doi: 10.1034/j.1399-0039.2003.00059.x.
38. Tsiavou A, Hatziagelaki E, Chaidaroglou A, Manginas A, Koniavitu K. TNF-alpha, TGF-beta1, IL-10, IL-6, gene polymorphisms in latent autoimmune diabetes of adults (LADA) and type 2 diabetes mellitus. *J Clin Immunol*. 2004;24(6):591-9.
39. Vendrell J, Fernandez-Real JM, Gutierrez C, Zamora A, Simon I. A polymorphism in the promoter of the tumor necrosis factor-alpha gene (2308) is associated with coronary heart disease in type 2 diabetic patients. *Atherosclerosis*. 2003;167(2):257-64. doi: 10.1016/S0021-9150(02)00429-X.
40. Dutta D, Choudhuri S, Mondal SA, Maisnam I. Tumor necrosis factor alpha -238G/A (rs 361525) gene polymorphism predicts progression to type-2 diabetes in an Eastern Indian population with prediabetes. *Diabetes Res Clin Pract*. 2013;99(3):e37-41. doi: 10.1016/j.diabres.2012.12.007.
41. Feng RN, Zhao C, Sun CH, Li Y. Meta-analysis of TNF 308 G/A polymorphism and type 2 diabetes mellitus. *PLoS one*. 2011;6(4):e18480. doi: 10.1371/journal.pone.0018480.
42. Feng RN, Li Y, Sun CH. TNF 308 G/A polymorphism and type 1 diabetes: a meta-analysis. *Diabetes Res Clin Pract*. 2009; 85(1):e4-7. doi: 10.1016/j.diabres.2009.04.026.
43. Mukhopadhyaya PN, Acharya A, Chavan Y, Purohit SS, Mutha A. Meta genomic study of single-nucleotide polymorphism within candidate genes massociated with type 2 diabetes in an Indian population. *Genet Mol Res*. 2010;9(4):2060-8.
44. Costa A, Fernandez-Real JM, Vendrell J, Broch M,

- Casamitjana R, Ricart W. Lower rate of tumor necrosis factor- $\alpha$  863a allele and higher concentration of tumor necrosis factor- $\alpha$  receptor 2 in first-degree relatives of subjects with type 2 diabetes. *Metabolism*. 2003;52(8):1068-71. doi: 10.1016/S0026-0495(03)00108-2.
45. Sookoian SC, Gonzalez C, Pirola CJ. Meta-analysis on the G-308A tumor necrosis factor alpha gene variant and phenotypes associated with the metabolic syndrome. *Obes Res*. 2005;13(12):2122-31. doi: 10.1038/oby.2005.263
  46. Liu ZH, Ding YL, Xiu LC, Pan HY. A meta-analysis of the association between TNF- $\alpha$  -308G>A polymorphism and type 2 diabetes mellitus in Han Chinese population. *PLoS One*. 2013;8(3):e59421. doi: 10.1371/journal.pone.0059421
  47. Bennett PH. Type 2 Diabetes: Prediction and Prevention. *Diabetic Med*. 2000;17(8):625-625. doi: 10.1046/j.1464-5491.2000.00290.x.
  48. Wu Y, Ding Y, Tanaka Y, Zhang W. Risk factors contributing to type 2 diabetes and recent advances in the treatment and prevention. *Int J Med Sci*. 2014;11(11):1185. doi: 10.7150/ijms.10001.
  49. Lillioja S, Mott DM, Spraul M, Ferraro R, Foley JE, Ravussin E. Insulin resistance and insulin secretory dysfunction as precursors of non-insulin-dependent diabetes mellitus. Prospective studies of Pima Indians. *N Engl J Med*. 1993;329(27):1988-92. doi: 10.1056/NEJM199312303292703.

**Copyright** © 2018 The Author(s); Published by Society of Diabetic Nephropathy Prevention. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.